

THE BRAUER GROUP OF A CURVE OVER A STRICTLY LOCAL DISCRETE VALUATION RING

TIMOTHY J. FORD

In memory of Professor S. A. Amitsur

ABSTRACT. Let K be the field of fractions of a curve over R where R is the henselization of a regular local ring on an algebraic curve over a field which is algebraically closed and has characteristic 0. Then K has the exponent = degree property for division algebras. In fact every central finite dimensional K -division algebra with exponent n is a cyclic algebra of degree n .

In this paper we continue to investigate the structure of division algebras D finite dimensional over their center K . The motivating problem is to classify those fields K that have the *exponent = degree* property for division algebras. We say that K has the *exponent = degree* property if for any central K -division algebra D the exponent of the class $[D]$ in the Brauer group $B(K)$ is equal to the degree $\sqrt{(D : K)}$ of the division algebra. Throughout this paper k is an algebraically closed field of characteristic 0.

Example 1. Some fields that are known to have the exponent = degree property are listed below.

- (1) A global field (an algebraic number field or a function field finitely generated of transcendence degree 1 over a finite field). This is classical.
- (2) The quotient field of either (a) the henselization $\mathcal{O}_{p,X}^h$ or (b) the completion $\hat{\mathcal{O}}_{p,X}$ at a closed point p on a normal surface X over k [1] or [6].
- (3) The quotient field of a ring obtained by (a) henselizing or (b) completing an affine surface over k along an integral curve [4].

In fact in each of these three examples, each division algebra D is split by a cyclic extension $K(\alpha^{1/n})$ for some $\alpha \in K$ and $n = \text{exponent}(D)$.

The purpose of this paper is to add to the list of Example 1 another class of fields satisfying the exponent = degree property.

Let $\mathcal{O}_{p,X}$ be the local ring at a regular point p on an algebraic curve X over the field k . Then $\mathcal{O}_{p,X}$ is a local principal ideal domain, hence a discrete valuation ring. The residue field of $\mathcal{O}_{p,X}$ is k . Let $R = \mathcal{O}_{p,X}^h$ be the henselization of $\mathcal{O}_{p,X}$. Consider an affine algebraic curve C over R . Following [9], C is an affine scheme together with a structure morphism $\pi : C \rightarrow \text{Spec } R$ such that π is flat and of finite type, the fibers of π are algebraic curves, and C is connected. Then π has 2 fibers. The closed fiber $\pi : C_0 \rightarrow x_0$ over the closed point x_0 of $\text{Spec } R$ is an algebraic curve over k . The open fiber $\pi : C_\eta \rightarrow \eta$ over the open point of $\text{Spec } R$ is an algebraic curve over the quotient field of R . Assume that C_η is integral, with

$K = K(C_\eta)$ the field of fractions. Our main result is that K has the exponent = degree property for division algebras.

Theorem 2. *Let R , C and K be as above and let D be a central finite dimensional K -division algebra with $\text{exponent}(D) = n$. Then D is a cyclic algebra of degree n .*

Proof. The proof is in the flavor of those used by [6] and [4].

Since C is $\text{Spec } S$ for an algebra S of finite type over R , we can assume C is a closed subscheme of affine space \mathbb{A}_R^m over R . Without changing K we can replace C with a projective completion over R . If necessary, we can also desingularize C . Therefore assume that $\pi : C \rightarrow \text{Spec } R$ is proper, that the open fiber is a nonsingular integral curve C_η over the quotient field of R . By Embedded Resolution of Curves in Surfaces [8, p. 391], we can assume that the closed fiber $(C_0)_{\text{red}}$ is a divisor over k with normal crossings. That is, write the reduced closed fiber $(C_0)_{\text{red}}$ as a union $C_1 \cup C_2 \cup \cdots \cup C_s$ of irreducible curves. By the normal crossing hypothesis we assume each component C_j is a nonsingular curve and that $(C_0)_{\text{red}}$ has at most ordinary double points as singularities.

Let L/K be a finite extension of fields and $Y \rightarrow C$ the integral closure of C in L . Let $f : Y' \rightarrow Y$ be any desingularization of Y . That is, Y' is nonsingular and f is a proper birational morphism. There is a complex

$$(1) \quad 0 \rightarrow B(Y') \rightarrow B(L) \xrightarrow{a} \bigoplus_{\Delta} H^1(K(\Delta), \mathbb{Q}/\mathbb{Z}) \xrightarrow{r} \bigoplus_P \mu(-1) \xrightarrow{s} H^4(Y', \mu) \rightarrow 0$$

which is exact except possibly at the term $\bigoplus H^1(K(\Delta), \mathbb{Q}/\mathbb{Z})$. The first summation is over all irreducible curves $\Delta \subseteq Y'$, the second over all closed points $P \in Y'$. This follows by combining sequences (3.1) and (3.2) of [2]. If $H^3(Y', \mu) = 0$, (1) is exact. The first two groups in (1) are the Brauer groups respectively, of Y' and L . The map a “measures the ramification” of a division algebra Λ over L . The ramification divisor of Λ is the set of divisors Δ where $a[\Lambda]$ is nontrivial. The group $H^1(K(\Delta), \mathbb{Q}/\mathbb{Z})$ classifies the cyclic Galois extensions of $K(\Delta)$ the function field of Δ . The map r measures the ramification of cyclic extensions of $K(\Delta)$. Here $\mu(-1) = \bigcup_n \text{Hom}(\mu_n, \mathbb{Q}/\mathbb{Z})$. Let D be a central K -division algebra

and $D_L = D \otimes L$, the restriction of D to L . We say that L splits the ramification of D on C if there exists a desingularization $f : Y' \rightarrow Y$ such that the class of D_L in the Brauer group $B(L)$ is in the image of the Brauer group $B(Y')$ of Y' .

We proceed as in the proof of [4, Cor. 5]. Since R is a direct limit of étale neighborhoods of (X, p) , C is of finite type over R and D is a finite K -algebra, we can find an étale neighborhood (U, p) of (X, p) , and a nonsingular algebraic surface C_1 satisfying the following.

- (1) There is a proper morphism $C_1 \rightarrow U$.
- (2) $C = C_1 \times_U \text{Spec } R$
- (3) If K_1 is the function field of C_1 , then there is a central simple algebra D_1 over K_1 such that $D = D_1 \otimes_{K_1} K$.

It was shown in the text immediately preceding Theorem 1.6 of [6] and again in [4, Prop. 3] that there exists a surface C_2 and a proper birational morphism $C_2 \rightarrow C_1$ and a cyclic field extension L_1/K_1 of degree n such that L_1 splits the ramification of D_1 on C_2 . Furthermore, if Y_1 is the integral closure of C_2 in L_1 , then Y_1 has only rational singularities. Let L denote the field KL_1 . Then L/K is

for each x in $\text{Spec } R$. So f is quasi-finite. Since f is a composite of proper morphisms, f is proper. But a proper quasi-finite morphism is finite [10, p. 6] so D_j is finite over R . Since R is henselian, any connected component of $D_j \times_R x_0$ gives rise to a connected component of D_j . But D_j is integral, hence $D_j \times_R x_0 = P_0$. So ${}_n B(C) = 0$ for each $n > 1$. \square

Lemma 4. *Let C be an algebraic curve over k embedded with normal crossings on a surface S . If the irreducible components of C are C_1, \dots, C_s , then $H^2(C, \mu_n) \cong \coprod_{i=1}^s H^2(C_i, \mu_n)$.*

Proof. Let $D = C_1 \coprod \dots \coprod C_s$ denote the disjoint union of the curves C_1, \dots, C_s . There is an obvious finite projection $\pi : D \rightarrow C$. Let σC denote the singular locus of C and $\sigma D = \pi^{-1}(\sigma C)$ those points on D lying over σC . Since C has only nodal singularities, the map $\pi : \sigma D \rightarrow \sigma C$ is 2-to-1. Let P be an element of σC and consider the cohomology with supports in P , $H_P^2(C, \mu_n)$. The strictly local ring $\mathcal{O}_{C,P}^h$ is henselian with algebraically closed residue field k . Let U denote $\text{Spec } \mathcal{O}_{C,P}^h$ and let P^h denote the closed point of U . By excision [10, Cor. 1.28, p. 93] $H_P^2(C, \mu_n) \cong H_{P^h}^2(U, \mu_n)$. The long exact sequence for $P^h \subseteq U$ is

$$(5) \quad \dots \rightarrow H^1(U, \mu_n) \rightarrow H^1(U - P^h, \mu_n) \rightarrow H_{P^h}^2(U, \mu_n) \rightarrow H^2(U, \mu_n) \rightarrow \dots$$

The curve U consists of 2 nonsingular henselian curves U_1, U_2 crossing at the closed point P^h . Each curve U_i is the prime spectrum of a henselian discrete valuation ring with residue field k . So $H^i(U, \mu_n) = 0$ for $i > 0$ and $H^1(U - P^h, \mu_n) = H^1(U_1 - P^h, \mu_n) \oplus H^1(U_2 - P^h, \mu_n) \cong \mathbb{Z}/n \oplus \mathbb{Z}/n$. So equation (5) and excision show that $H_P^2(C, \mu_n) \cong (\mathbb{Z}/n)^{(2)}$. If Q is an element of σD , then the argument above also shows that $H_Q^2(D, \mu_n) \cong \mathbb{Z}/n$. Since σC decomposes into a finite number of points P , it follows that $H_{\sigma C}^2(C, \mu_n)$ decomposes into the direct sum $\coprod_{P \in \sigma C} H_P^2(C, \mu_n)$ and similarly for σD . The long exact sequences of cohomology with supports in σC and σD combined with the maps induced by π yield the commutative diagram below.

$$(6) \quad \begin{array}{ccccccc} H^1(C - \sigma C, \mu_n) & \longrightarrow & H_{\sigma C}^2(C, \mu_n) & \longrightarrow & H^2(C, \mu_n) & \longrightarrow & H^2(C - \sigma C, \mu_n) \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \delta \\ H^1(D - \sigma D, \mu_n) & \longrightarrow & H_{\sigma D}^2(D, \mu_n) & \longrightarrow & H^2(D, \mu_n) & \longrightarrow & H^2(D - \sigma D, \mu_n) \end{array}$$

Because $\pi : C - \sigma C \xrightarrow{\cong} D - \sigma D$, α and δ are isomorphisms. The map β is an isomorphism by the above computations. Therefore γ is an isomorphism. Because D is a disjoint union, the lemma follows. \square

Corollary 5. *Let R, C and K be as in Theorem 3. The sequence*

$$0 \rightarrow B(K) \xrightarrow{a} \bigoplus_{\Delta} H^1(K(\Delta), \mathbb{Q}/\mathbb{Z}) \xrightarrow{r} \bigoplus_P \mu(-1) \rightarrow 0$$

is exact where the first summation is over all irreducible curves $\Delta \subseteq C$, the second over all closed points $P \in C$.

Proof. Follows immediately from (1) and Theorem 3. \square

For any discrete valuation ring R with perfect residue field k and field of fractions K , for each $q \geq 0$ the natural map $H^q(R, \mathbf{G}_m) \rightarrow H^q(k, \mathbf{G}_m)$ is an isomorphism [7, III, p. 93]. For $q = 2$ this is the theorem of Azumaya. There is a split-exact sequence

$$0 \rightarrow H^q(k, \mathbf{G}_m) \rightarrow H^q(K, \mathbf{G}_m) \rightarrow H^{q-1}(k, \mathbf{G}_m) \rightarrow 0$$

for each $q \geq 2$ [7, III, p. 93 and p. 188]. In particular, for $q = 2$, it follows that the Brauer group of K decomposes into a direct sum of $B(k)$ and $H^1(k, \mathbf{Q}/\mathbf{Z})$. The group $H^1(k, \mathbf{Q}/\mathbf{Z})$ parametrizes the unramified cyclic Galois extensions of K . Various other results along these same lines are derived in [3] and [11]. We arrive at similar results for the Brauer group of a curve over K .

Theorem 6. *Let R be a strictly local ring which is the henselization of a local ring $\mathcal{O}_{p,X}$ at a closed point p on a smooth curve X over k . Let $\pi : C \rightarrow \text{Spec } R$ be proper and smooth of relative dimension 1. Let η be the generic point of $\text{Spec } R$ and $C_\eta = C \times_R \eta$. Let x_0 be the closed point of $\text{Spec } R$ and $C_0 = C \times_R x_0$. Then the Brauer group $B(C_\eta)$ is isomorphic to $H^1(C_0, \mathbf{Q}/\mathbf{Z})$. Every Azumaya algebra over C_η of exponent n is split by a cyclic Galois extension of degree n which descends to an unramified extension of C .*

Proof. Since $\text{Spec } R = \{x_0\} \cup \{\eta\}$ with $\{x_0\}$ closed and $\{\eta\}$ open, we have $C = C_0 \cup C_\eta$ with C_0 closed and C_η open. The long exact sequence of cohomology with supports in C_0 and coefficients in \mathbf{G}_m is

$$(7) \quad \begin{aligned} H^0(C, \mathbf{G}_m) &\rightarrow H^0(C_\eta, \mathbf{G}_m) \rightarrow H^1_{C_0}(C, \mathbf{G}_m) \rightarrow \\ &H^1(C, \mathbf{G}_m) \rightarrow H^1(C_\eta, \mathbf{G}_m) \rightarrow H^2_{C_0}(C, \mathbf{G}_m) \rightarrow \\ &H^2(C, \mathbf{G}_m) \rightarrow H^2(C_\eta, \mathbf{G}_m) \rightarrow H^3_{C_0}(C, \mathbf{G}_m) \rightarrow \dots \end{aligned}$$

If t is a local parameter for R , then $\frac{H^0(C_\eta, \mathbf{G}_m)}{H^0(C, \mathbf{G}_m)} = \langle t \rangle$ and t vanishes with order 1 along C_0 . Since C_0 is a principal divisor, $\text{Pic } C \cong \text{Pic } C_\eta$. From Kummer theory the diagram

$$(8) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H^0(C_\eta, \mathbf{G}_m) \otimes \mathbf{Z}/n & \longrightarrow & H^1(C_\eta, \mu_n) & \longrightarrow & {}_n\text{Pic } C_\eta \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \cong \\ 0 & \longrightarrow & H^0(C, \mathbf{G}_m) \otimes \mathbf{Z}/n & \longrightarrow & H^1(C, \mu_n) & \longrightarrow & {}_n\text{Pic } C \longrightarrow 0 \end{array}$$

commutes and has exact rows. Since $H^0(C_\eta, \mathbf{G}_m) \otimes \mathbf{Z}/n \cong \langle t \rangle / \langle t^n \rangle$ and $H^0(C, \mathbf{G}_m) \otimes \mathbf{Z}/n = 0$, we see from (8) that $H^1(C_\eta, \mu_n) \cong \langle t \rangle / \langle t^n \rangle \times H^1(C, \mu_n)$ and by proper base change $H^1(C, \mu_n) \cong H^1(C_0, \mu_n)$. From Theorem 3 we have $H^q(C, \mathbf{G}_m) = 0$ for all $q \geq 2$. From (7) it follows that $H^q(C_\eta, \mathbf{G}_m) \cong H^q_{C_0}(C, \mathbf{G}_m)$ for all $q \geq 2$. Since $\text{Spec } R$ is a direct limit of étale neighborhoods $U \hookrightarrow X$ of the closed point $p \hookrightarrow X$, the morphism $\pi : C \rightarrow \text{Spec } R$ descends to a proper smooth morphism $C' \rightarrow U$ with a closed fiber $C'_0 = C' \times_U p \cong C_0$. By cohomological purity (see [5]) $H^q_{C'_0}(C', \mathbf{G}_m) \cong H^{q-2}(C_0, \mathbf{Q}/\mathbf{Z})$ for all $q \geq 3$. Taking the limit over all such $U \hookrightarrow X$ yields $H^q_{C_0}(C, \mathbf{G}_m) \cong H^{q-2}(C_0, \mathbf{Q}/\mathbf{Z})$ for all $q \geq 3$. Therefore $H^2(C_\eta, \mathbf{G}_m) \cong H^1(C_0, \mathbf{Q}/\mathbf{Z})$, $H^3(C_\eta, \mathbf{G}_m) \cong H^2(C_0, \mathbf{Q}/\mathbf{Z}) \cong \mu$ and $H^q(C_\eta, \mathbf{G}_m) \cong H^{q-1}(C_0, \mathbf{Q}/\mathbf{Z}) = 0$ for all $q \geq 4$. The isomorphism ${}_n H^2(C_\eta, \mathbf{G}_m) \cong H^1(C_0, \mathbf{Z}/n)$

is the Gysin map [10, p. 244]. Given any λ in $H^1(C, \mathbb{Z}/n)$ there is a corresponding λ_0 in $H^1(C_0, \mathbb{Z}/n)$. Let λ_η denote the image of λ in the group $H^1(C_\eta, \mathbb{Z}/n)$ under the natural map. Let λ_η also denote the corresponding cyclic Galois cover of C_η with group $\langle \sigma \rangle$. Using the cyclic Galois cover λ_η and the trivial factor set t we form a cyclic crossed product algebra $\Delta(\lambda) = (\lambda_\eta/C_\eta, \sigma, t)$ which represents a class in ${}_n\mathbb{B}(C_\eta)$. Consider the ramification divisor of $\Delta(\lambda)$ on C . Along the divisor C_0 of C , the ramification of $\Delta(\lambda)$ is the element λ_0 of the group $H^1(C_0, \mathbb{Z}/n)$. Therefore, the correspondence $\lambda \mapsto \Delta(\lambda)$ induces an isomorphism $H^1(C, \mathbb{Z}/n) \cong {}_n\mathbb{B}(C_\eta)$. Every Azumaya algebra over C_η whose Brauer class is annihilated by n is Brauer equivalent to a cyclic crossed product of the form $(\lambda_\eta/C_\eta, \sigma, t)$, hence is split by λ_η for some λ . \square

REFERENCES

- [1] M. Artin, *Two dimensional orders of finite representation type*, Manuscripta Math., **58** (1987), pp. 445–471.
- [2] M. Artin and D. Mumford, *Some elementary examples of unirational varieties which are not rational*, Proc. London Math. Soc. (3), **25** (1972), pp. 75–95.
- [3] M. Auslander and A. Brumer, *Brauer groups of discrete valuation rings*, Nederl. Akad. Wetensch. Proc. Ser. A, **71** (1968), pp. 286–296.
- [4] T. J. Ford, *Division algebras over nonlocal henselian surfaces*, Pacific J. Math., **147** (1991), pp. 301–310.
- [5] ———, *On the Brauer group of a localization*, J. Algebra, **147** (1992), pp. 365–378.
- [6] T. J. Ford and D. Saltman, *Division algebras over henselian surfaces*, Israel Mathematical Conference Proceedings, **1** (1989), pp. 320–336.
- [7] A. Grothendieck, *Le groupe de Brauer I, II, III*, in Dix Exposés sur la Cohomologie des Schémas, North Holland, Amsterdam, 1968, pp. 46–188.
- [8] R. Hartshorne, *Algebraic Geometry*, vol. 52 of Graduate Texts in Mathematics, Springer-Verlag, New York/Berlin, 1977.
- [9] S. Lichtenbaum, *Curves over discrete valuation rings*, Amer. J. Math., **90** (1968), pp. 380–405.
- [10] J. Milne, *Etale Cohomology*, vol. 33 of Princeton Mathematical Series, Princeton University Press, Princeton, N.J., 1980.
- [11] S. Yuan, *On the Brauer groups of local fields*, Ann. of Math. (2), **82** (1965), pp. 434–444.

DEPARTMENT OF MATHEMATICS, FLORIDA ATLANTIC UNIVERSITY, BOCA RATON, FL 33431
E-mail address: Ford@acc.fau.edu